

HUB EP-98/43
DFPD 98/TH/36
hep-th/9807133

Harmonics, Notophs and Chiral Bosons

Paolo Pasti*, Dmitri Sorokin *, **¹ and Mario Tonin*

* Università Degli Studi Di Padova, Dipartimento Di Fisica “Galileo Galilei”
ed INFN, Sezione Di Padova
Via F. Marzolo, 8, 35131 Padova, Italia

** Humboldt-Universität zu Berlin
Mathematisch-Naturwissenschaftliche Fakultät
Institut für Physik
Invalidenstrasse 110, D-10115 Berlin, Germany

ABSTRACT

A way of covariantizing duality symmetric actions is described.

The presence of self-dual fields or, in more general case, duality-symmetric fields in field-theoretical and string models reflects their duality properties whose extreme importance for understanding a full quantum theory has been appreciated during an impetuous development of the duality field happened during last few years. The knowledge of duality-symmetric effective actions is useful for carrying out more systematic study of the classical and quantum properties of the theory, and in this memorial contribution we would like to demonstrate how fruitful physical ideas and mathematical techniques which Victor Isakovich Ogievetsky and his colleagues have developed helped us to construct a covariant Lagrangian formulation applicable to all known models with duality-symmetric fields in space-time of Lorentz signature.

The problem of constructing and studying models described by duality-invariant actions has a rather long history. It goes back to time when Poincare and later on Dirac noticed electric-magnetic duality symmetry of the free Maxwell equations, and, Dirac assumed the existence of magnetically charged particles (monopoles and dyons) [1] admitting the duality symmetry to be also held for the Maxwell equations in the presence of charged sources. To describe monopoles and dyons on an equal footing with electrically charged particles one should have a duality-symmetric form of the Maxwell action. This problem was studied (among others) by Schwinger and

¹ Alexander von Humboldt fellow.

On leave from Kharkov Institute of Physics and Technology, Kharkov, 310108, Ukraine.

Zwanziger, and in 1971 Zwanziger proposed a duality-symmetric action for Maxwell fields interacting with dyonic sources [2]. An alternative duality-symmetric Maxwell action was proposed by Deser and Teitelboim in 1976 [4]. The two actions, which proved to be dual to each other [3], are not manifestly Lorentz-invariant. This feature turned out to be a general one. Duality and space-time symmetries hardly coexist in one and the same action.

Later on this problem arose in multidimensional supergravity theories in space-time of a dimension $D = 4p + 2$ where one would like to know how to construct an action for self-dual tensor fields (chiral bosons) which are present in some versions of supergravity and in the heterotic string. One of the ways of solving this problem is to sacrifice Lorentz covariance in favour of duality symmetry. A non-covariant action for $D = 2$ chiral bosons was constructed by Floreanini and Jackiw [5], and Henneaux and Teitelboim [6] proposed non-covariant actions for self-dual fields in higher dimensional $D = 4p + 2$ space-time. In a context of modern aspects of duality Tseytlin [7] considered a duality-symmetric action for a string. Finally, Schwarz and Sen [8] constructed non-covariant duality-symmetric actions for dual tensor fields in any space-time dimension.

There have also been developed covariant approaches to the construction of duality-symmetric actions. These use auxiliary fields. The first covariant Lagrangian formulation of chiral bosons was proposed by Siegel [9] and its modification was considered by Kavalov and Mkrtchyan [10] in application to D=6 and D=10 chiral supergravities. Another covariant approach is based on the use of an infinite number of auxiliary fields [11, 12]. It might be interesting that an effective self-dual action of this kind was extracted from a type IIB string field theory [12].

The third formulation was proposed in [13]. In its minimal version only one scalar auxiliary field is used to ensure space-time covariance of duality-symmetric actions. This approach turned out to be the most appropriate for the construction of the worldvolume action for the M-theory five-brane [14], duality-symmetric D=11 supergravity [15] and D=10 IIB supergravity [16].

Below we will use Maxwell theory to demonstrate how this third approach was developed with promptings provided by works of V. I. Ogievetsky.

It is well known that the standard action for a free Maxwell field is not invariant under duality transformations of its electric and magnetic strength vectors. To have a duality symmetry at the level of action one should double the number of gauge fields (A_m^α , $\alpha = 1, 2$, $m=0,1,2,3$) [2, 4, 8] and construct an action in such a way that one of the gauge fields becomes an auxiliary field upon solving equations of motion. The duality symmetric action of refs. [4, 8] can be written in the following form:

$$S = \int d^4x \left(-\frac{1}{8} F_{mn}^\alpha F^{mn\alpha} + \frac{1}{4} \mathcal{F}_{0i}^\alpha \mathcal{F}_{i0}^\alpha \right), \quad (i = 1, 2, 3) \quad (1)$$

where

$$\mathcal{F}_{mn}^\alpha = \mathcal{L}^{\alpha\beta} F_{mn}^\beta - \frac{1}{2} \epsilon_{mnlp} F^{lp\alpha} = \frac{1}{2} \epsilon_{mnlp} \mathcal{F}^{lp\beta} \mathcal{L}^{\alpha\beta}, \quad (2)$$

($\mathcal{L}^{12} = -\mathcal{L}^{21} = 1$) is the self-dual combination of the field strengths.

The Zwanziger action [2] differs from (1) by the sign in front of the second term and in that, instead of the time-coordinate index, one of the spatial indices is separated in the analogous term of the Zwanziger action.

Duality symmetry is a discrete subgroup of $SO(2)$ rotations of A_m^α ($A_m^\alpha \rightarrow \mathcal{L}^{\alpha\beta} A_m^\beta$).

Note that because of the self-duality property (2) $\mathcal{F}_{mn}^\alpha \mathcal{F}^{\alpha mn} \equiv 0$, and the best thing which one can do is to take the square of only a part of the components of $\mathcal{F}_{mn}^\alpha$ for the construction of the second term of the action (1), and this breaks manifest Lorentz invariance.

Here is a place to explain why the signature of space-time is important for the possibility of applying the Lagrangian approach considered to the description of chiral bosons. It is crucial for this approach that the “square” of a self-dual tensor is zero, which holds, for instance, in $D = 2p+2$ spaces of a Lorentz signature. Then taking the square of an appropriate part of the components of the self-dual tensor (as in (1)) one gets the desirable result. On the contrary, for instance, in $D = 4$ space of Euclidian signature the square of the self-dual combination of a gauge field-strength is non-zero and reproduces (up to a total derivative) the standard Maxwell Lagrangian, and no reasonable choice of its components is known in these cases to construct actions analogous to (1).

We have seen that the method we used to get the action breaks manifest Lorentz invariance, however, beside the manifest spatial rotations the action (1) is invariant under the following modified space-time transformations of A_i^α (in the gauge $A_0^\alpha = 0$)

$$\delta A_i^\alpha = x^0 v^k \partial_k A_i^\alpha + v^k x^k \partial_0 A_i^\alpha + v^k x^k \mathcal{L}^{\alpha\beta} \mathcal{F}_{0i}^\beta, \quad (3)$$

where the first two terms describe the ordinary Lorentz boosts along a constant velocity v^i and the third term vanishes on the mass shell since an additional local symmetry of the action (1)

$$\delta A_0^\alpha = \varphi^\alpha(x) \quad (4)$$

allows one to reduce the equations of motion

$$\frac{\delta S}{\delta A_i^\alpha} = \epsilon^{ijk} \partial_i \mathcal{F}_{k0}^\alpha = 0 \quad (5)$$

to the duality condition

$$\mathcal{F}_{mn}^\alpha = \mathcal{L}^{\alpha\beta} F_{mn}^\beta - \frac{1}{2} \epsilon_{mnlp} F^{lp\alpha} = 0 \quad (6)$$

which, on the one hand, leads to the Maxwell equations

$$\partial_m \mathcal{F}^{mn\alpha} = \partial_m F^{mn\alpha} = 0 \quad (7)$$

and, on the other hand, completely determines one of the gauge fields through another one. For instance, using the relation

$$\frac{1}{2} \epsilon_{mnlp} F^{2mn} = F_{mn}^1$$

we can exclude $A_m^2(x)$ from the action (1) and get the conventional Maxwell action.

One can admit that the action (1) arose as a result of some gauge fixing which specifies time direction in a Lorentz covariant action [13].

The first step is to covariantize the self-dual part of the action (1). For people who are acquainted with harmonic techniques served for similar covariantization purposes in supersymmetric theories [17] the first thing which comes into mind is to introduce an auxiliary harmonic-like vector field

$$l_m(x) \equiv \frac{u_m(x)}{\sqrt{-u_n u^n}}, \quad l_m l^m = -1, \quad (8)$$

and to write the action as follows:

$$S_A = \int d^4x \left(-\frac{1}{8} F_{mn}^\alpha F^{\alpha mn} + \frac{1}{4(-u_l u^l)} u^m \mathcal{F}_{mn}^\alpha \mathcal{F}^{\alpha np} u_p \right). \quad (9)$$

The field (8) is harmonic in the sense that if supplemented with space-like fields $l_m^i(x)$, $l_m^i l^{jm} = \delta^{ij}$, $l^m l_m^i = 0$ (which do not enter the action) the set of the four vector fields form a matrix of the Lorentz group $SO(1, 3)$ and can be used to contract Lorentz indices of other fields in a covariant way. For the analysis of properties of the action it was proved more convenient to work with the field $u(x)$ rather than with its normalized form $l(x)$, and at the same time to use harmonic properties of the latter.

The main problem is to find a local symmetry which would permit to choose a gauge where u_m is a constant vector, in particular,

$$u_m(x) = \delta_m^0. \quad (10)$$

Then the action (9) can reduce to (1). Note that a spatial gauge, for instance $u_m(x) = \delta_m^3$ is equally admissible and leads to a non-covariant action which also produces the duality condition (6).

The search for this symmetry turns out to be connected with another problem, namely, the problem of preserving a local symmetry under (4). In the covariant version this transformation should be replaced by

$$\delta A_m^\alpha = u_m \varphi^\alpha. \quad (11)$$

To keep this symmetry is important (as we have already seen) for getting the duality condition (6).

To have the invariance under transformations (11) one should add to the Lorentz invariant action (9) another term

$$S_B = - \int d^4x \epsilon^{mnpq} u_m \partial_n B_{pq}, \quad (12)$$

where $B_{mn}(x)$ is an antisymmetric tensor field. Then the variation of (9) under (11) is canceled by the variation of (12) under

$$\delta B_{mn} = -\frac{\varphi^\alpha}{u^2} (\mathcal{F}_{mp}^\alpha u^p u_n - \mathcal{F}_{np}^\alpha u^p u_m). \quad (13)$$

The equation of motion of B_{mn}

$$\partial_{[m}u_{n]} = 0 \quad \rightarrow \quad u_m(x) = \partial_m a(x) \quad (14)$$

reads that $u_m(x)$ is the derivative of a scalar field. Note also that (12) is invariant under

$$\delta B_{mn} = \partial_{[m}b_{n]}(x). \quad (15)$$

As in the case of the action (1), the local symmetry (11) allows one to fix a gauge on the mass shell in such a way that the duality condition (6) takes place. To arrive at eq. (6) harmonic techniques found to be rather useful. Let us sketch the derivation of (6).

The equation of motion of A_m^α produced by (9) is

$$\epsilon^{mnpq}\partial_l(l_p\mathcal{F}_q^\alpha) = 0, \quad (16)$$

where $\mathcal{F}_q^\alpha \equiv \mathcal{F}_{qp}^\alpha l^p$ and l_p is defined by (8) and (14).

From (16) it follows that

$$l_{[p}\mathcal{F}_{q]}^\alpha = \partial_{[p}\Phi_{q]}^\alpha, \quad \mathcal{F}_q^\alpha = l^p\partial_p\Phi_q^\alpha - l^p\partial_q\Phi_p^\alpha, \quad (17)$$

where Φ_q^α are two vector functions. Projecting (17) onto harmonics l_i^p ($i = 1, 2, 3$) orthogonal to l_p we get

$$l_{[i}^p l_{j]}^q \partial_p \Phi_q^\alpha = 0,$$

which implies that

$$l_i^q \Phi_q^\alpha = l_i^q \partial_q \varphi^\alpha(x) \quad \Rightarrow \quad \Phi_q^\alpha = l_q \Phi^\alpha + \partial_q \varphi^\alpha(x). \quad (18)$$

Substituting (18) into (17) and taking into account that the last term in (18) can be neglected since (17) is invariant under gauge transformations $\Phi_q^\alpha \rightarrow \Phi_q^\alpha + \partial_q \varphi^\alpha(x)$ we obtain

$$\mathcal{F}_q^\alpha = \partial_q \Phi^\alpha(x) + l^p \partial_p (l_q \Phi^\alpha). \quad (19)$$

Now the transformation (11) can be used to put in (19) $\Phi^\alpha = 0$ as a gauge fixing condition. Then we have $\mathcal{F}_q^\alpha = 0$ which, because of the self-duality of $\mathcal{F}_{pq}^\alpha$, implies (6).

Thus, we again remain with only one independent Maxwell field and get the duality between its electric and magnetic strength vector. In view of the vanishing condition for the self-dual strength tensor the equations of motion of u_m reduce to:

$$\frac{\delta(S_A + S_B)}{\delta u_m} = \epsilon^{mnlp} \partial_n B_{lp} = 0 \quad \rightarrow \quad B_{mn} = \partial_{[m}b_{n]}, \quad (20)$$

which means that B_{mn} is completely auxiliary and can be eliminated by use of the corresponding local transformations (15).

The only thing which has remained to show is that u_m itself does not carry physical degrees of freedom and can be gauge fixed to $u_m = \delta_m^0$. For this we have to find a corresponding local symmetry. And here an analogy of the antisymmetric field B_{mn} with the ‘notoph’ of Ogievetsky and Polubarinov [18] helps us to get the corresponding symmetry transformations.

The form of the action (12) containing B_{mn} reminds a term which one encounters in a dual formulation of a pseudoscalar (‘axion’) field as an antisymmetric notoph field (see, for instance, [18, 19])

$$S = \int d^4x \left(-\frac{1}{2}(\partial_m a(x) - u_m(x))(\partial^m a(x) - u^m(x)) - \epsilon^{pqmn} u_p \partial_q B_{mn} \right). \quad (21)$$

The action (21) is invariant under local Peccei–Quinn transformations

$$\delta a(x) = \varphi(x), \quad \delta u_m = \partial_m \varphi(x), \quad (22)$$

(u_m being the corresponding gauge field) and produces dual versions of the axion action²:

$$\begin{aligned} L &= -\frac{1}{2} \partial_m a(x) \partial^m a(x), \\ L &= \frac{1}{3!} \partial_{[m} B_{np]} \partial^{[m} B^{np]}. \end{aligned} \quad (23)$$

The duality relation between the pseudoscalar field $a(x)$ and the antisymmetric tensor field B_{mn}

$$\partial_l a(x) = \epsilon_{lmnp} \partial^m B^{np} \quad (24)$$

is a consequence of the equations of motion of u_m obtained from (21).

Now one can assume that the action (9)+(12) is also invariant under the transformations (22). This is indeed the case provided A_m^α and B_{mn} transform as follows

$$\delta A_m^\alpha = \frac{\varphi(x)}{u^2} \mathcal{L}^{\alpha\beta} \mathcal{F}_{mn}^\beta u^n, \quad \delta B_{mn} = \frac{\varphi(x)}{(u^2)^2} \mathcal{F}_m^{\alpha r} u_r \mathcal{F}_n^{\beta s} u_s \mathcal{L}^{\alpha\beta}. \quad (25)$$

Then, taking into account (14) and requiring that $u^2 \neq 0$ (to escape singularities), we can use the local transformation (22) to put $u_m = \delta_m^0$. In this gauge the manifestly Lorentz invariant duality–symmetric action

$$S = \int d^4x \left(-\frac{1}{8} F_{mn}^\alpha F^{\alpha mn} - \frac{1}{4(u_l u^l)} u^m \mathcal{F}_{mn}^\alpha \mathcal{F}^{\alpha np} u_p - \epsilon^{mnpq} u_m \partial_n B_{pq} \right) \quad (26)$$

reduces to (1), and the local transformations of A_m^α (25) (with $\varphi(x) = x^i v^i$) are combined with the corresponding Lorentz transformations and produce the modified space–time symmetry (3) of the action (1).

²We denoted the scalar field in (14) with the same letter $a(x)$ as the axion field to point to their formal “generic roots”.

One can reduce the number of the auxiliary fields in the action (26) to one scalar field by substituting into (26) the solution of the equation of motion (14). The resulting minimal form of the covariant duality symmetric action

$$S = \int d^4x \left(-\frac{1}{8} F_{mn}^\alpha F^{\alpha mn} - \frac{1}{4(\partial_l a \partial^l a)} \partial^m a(x) \mathcal{F}_{mn}^\alpha \mathcal{F}^{\alpha np} \partial_p a(x) \right), \quad (27)$$

which remains the same in all space–time dimensions, has been used for the description of chiral bosons in various theoretical models (see [14, 15, 16] and references therein).

We have thus obtained a covariant Lagrangian formulation of Maxwell theory which is also invariant under electric–magnetic duality. The action was shown to produce in the temporal gauge (10) the duality–symmetric action (1) of [4, 8].

As has been mentioned in the introduction the first duality–symmetric action for the Maxwell fields was constructed by Zwanziger [2] and that it differs from (1) in the sign of the second term. This difference leads to essentially different symmetry properties of the Zwanziger action [3]. However, since both actions describe one and the same physical model there should be a relationship between them. This relation is established through a duality transform of the auxiliary scalar field $a(x)$ into the auxiliary ‘notoph’ field B_{mn} [3]. For this consider (26) as a master action which produces different dual actions depending on which auxiliary fields are integrated out. The action (27) is one of these dual actions. Another one is obtained by varying (26) with respect to u_m , which gives an expression for the dual field strength $v^m = \epsilon^{mnpq} \partial_n B_{pq}$ in terms of u_m and the Maxwell field strengths, solving this expression for the vector field u_m in terms of v_m and substituting the result back into the action (26). The action now contains only the dual field strength v_m of the auxiliary field B_{mn} and has the form

$$S = \int d^4x \left(-\frac{1}{8} F_{mn}^\alpha F^{\alpha mn} + \frac{1}{4(v_l v^l)} v^m \mathcal{F}_{mn}^\alpha \mathcal{F}^{\alpha np} v_p \right), \quad (28)$$

the sign of the second term being changed with respect to the analogous term in (26) and (27). It can be shown that the action (28) is invariant under local transformations which allow one to fix $v_m(x)$ to be a constant time–like or space–like vector upon which (28) reduces to the Zwanziger action (see [3] for details).

In conclusion we have shown how the covariant Lagrangian approach to the description of duality–symmetric fields unifies different non–covariant formulations. This approach is also related to the infinite field approach [11] being a consistent truncation of the latter (see the last ref. in [13]).

From the action (27) one can *formally* obtain the Siegel action [9] by replacing $\frac{\partial_m a(x) \partial_p a(x)}{(\partial_l a \partial^l a)}$ with a Lagrange multiplier field $\Lambda_{mp}(x)$. But this relation is only a formal one since “hiding” derivatives of fields in other fields is not an innocent trick. The properties of the two actions are very different, a main difference being that the duality condition (6) is obtained from the actions (1), (26)–(28) as a consequence

of equations of motion of the gauge fields $A_m^\alpha(x)$, while in the Siegel formulation it arises as a “square root” of the constraint produced by the Lagrange multiplier Λ_{mn} equation of motion. This results in a different structure of Hamiltonian constraints and, as a consequence, leads to different ways of quantizing chiral bosons. We also note that for more complicated cases of self-interacting chiral bosons, such as the M-theory five-brane [14] an effective action in the Siegel form is not known yet.

Acknowledgements. This work was partially supported by the European Commission TMR Programme ERBFMPX-CT96-0045 to which the authors are associated, and by the INTAS Grant 96-308.

References

1. P. A. M. Dirac, *Proc. R. Soc.* **A133**, 60 (1931); *Phys. Rev.* **74**, 817 (1948).
2. D. Zwanziger, *Phys. Rev.* **D3**, 880 (1971).
3. A. Maznytsia, C. R. Preitschopf and D. Sorokin, Duality of self-dual actions, hep-th/9805110.
4. S. Deser and C. Teitelboim, *Phys. Rev.* **D13**, 1592 (1976).
5. R. Floreanini and R. Jackiw, *Phys. Rev. Lett.* **59**, 1873 (1987).
6. M. Henneaux and C. Teitelboim, in *Proc. Quantum mechanics of fundamental systems 2*, Santiago, 1987, p. 79; *Phys. Lett.* **B206**, 650 (1988).
7. A. Tseytlin, *Phys. Lett.* **B242**, 163 (1990); *Nucl. Phys.* **B350**, 395 (1991).
8. J. H. Schwarz and A. Sen, *Nucl. Phys.* **B411**, 35 (1994).
9. W. Siegel, *Nucl. Phys.* **B238**, 307 (1984).
10. A. R. Kavalov and R. L. Mkrtychyan, *Nucl. Phys.* **B331**, 391 (1990).
11. B. McClain, Y. S. Wu and F. Yu, *Nucl. Phys.* **B343**, 689 (1990);
C. Wotzasek, *Phys. Rev. Lett.* **66**, 129 (1991);
I. Martin and A. Restuccia, *Phys. Lett.* **B323**, 311 (1994);
F.P. Devecchi and M. Henneaux, *Phys. Rev.* **D45**, 1606 (1996);
I. Bengtsson and A. Kleppe, *Int. J. Mod. Phys.* **A12**, 3397 (1997).
12. N. Berkovits, *Phys. Lett.* **B388**, 743 (1996).
13. P. Pasti, D. Sorokin and M. Tonin, *Phys. Lett.* **B352**, 59 (1995); *Phys. Rev.* **D52**, R4277 (1995); *Phys. Rev.* **D55**, 6292 (1997).
14. P. Pasti, D. Sorokin and M. Tonin, *Phys. Lett.* **398B**, 41 (1997);
I. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. Sorokin and M. Tonin, *Phys. Rev. Lett.* **78**, 4332 (1997);
M. Aganagic, J. Park, C. Popescu and J. H. Schwarz, *Nucl. Phys.* **B496**, 191 (1997).
15. I. Bandos, N. Berkovits and D. Sorokin, *Nucl. Phys.* **B522** (1998) 214.
16. G. Dall’Agata, K. Lechner and D. Sorokin, *Class. Quant. Grav.* **14**, L195 (1997);
G. Dall’Agata, K. Lechner and M. Tonin, D=10, N=IIB supergravity: Lorentz-invariant actions and duality, hep-th/9806140.

- 17. A. Galperin, E. Ivanov, S. Kalitzin, V. Ogievetsky and E. Sokatchev, *Class. Quantum Grav.* **1**, 498 (1984); *Class. Quantum Grav.* **2**, 155 (1985).
- 18. V. I. Ogievetsky and I. V. Polubarinov, *Soviet. J. Nucl. Phys.* **4**, 156 (1967);
M. Kalb and P. Ramond, *Phys. Rev.* **D9** 2273 (1974).
- 19. S Coleman, J. Preskill and F. Wilczek, *Nucl. Phys.* **B378**, 175 (1992).